

juvenile *B. glabrata* infected with *E. caproni* larvae showed reduced fecundity.

Literature Cited

- Christensen, N. Ø., F. Frandsen, and M. Z. Roushdy. 1980. The influence of environmental conditions and parasite-intermediate host-related factors on the transmission of *Echinostoma liei*. *Zeitschrift für Parasitenkunde* 63:47–63.
- Crews, A. P., and T. P. Yoshino. 1989. *Schistosoma mansoni*: effect of infection on reproduction and gonadal growth in *Biomphalaria glabrata*. *Experimental Parasitology* 68:326–334.
- Fried, B., and J. E. Huffman. 1996. The biology of the intestinal trematode *Echinostoma caproni*. *Advances in Parasitology* 38:311–368.
- Hurd, H. 1990. Physiological and behavioural interactions between parasites and invertebrate hosts. *Advances in Parasitology* 29:272–317.
- Kuris, A. M. 1980. Effect of exposure to *Echinostoma liei* miracidia on growth and survival of young *Biomphalaria glabrata* snails. *International Journal for Parasitology* 10:303–308.
- Mueleman, E. A. 1972. Host–parasite interrelationships between the freshwater pulmonate *Biomphalaria pfeifferi* and the trematode *Schistosoma mansoni*. *Netherlands Journal of Zoology* 22:355–427.
- Raymond, K., and A. J. Probert. 1993. The effect of infection with *Schistosoma margrebowiei* on the growth of *Bulinus natalensis*. *Journal of Helminthology* 67:10–16.
- Thompson, S. N. 1985. Review: metabolic integration during the host associations of multicellular animal endoparasites. *Comparative Biochemistry and Physiology* 81B:21–42.
- . 1997. Physiology and biochemistry of snail larval trematode relationships. Pages 149–195 in B. Fried and T. K. Graczyk, eds. *Advances in Trematode Biology*. CRC, Boca Raton, Florida.
- Ulmer, M. J. 1970. Notes on rearing of snails in the laboratory. Pages 143–144 in A. J. MacInnis and M. Voge, eds. *Experiments and Techniques in Parasitology*. W. H. Freeman, San Francisco.

J. Helminthol. Soc. Wash.
65(1), 1998 pp. 98–102

Research Note

Helminth Parasites of the Spotted Salamander *Ambystoma maculatum* and Red-backed Salamander *Plethodon c. cinereus* from Northwestern Wisconsin

MATTHEW G. BOLEK AND JAMES R. COGGINS

Department of Biological Sciences, University of Wisconsin–Milwaukee, Milwaukee, Wisconsin 53201
(e-mail: bolekm@csd.uwm.edu, coggins@csd.uwm.edu)

ABSTRACT: Twenty spotted salamanders *Ambystoma maculatum* and 20 red-backed salamanders *Plethodon c. cinereus* were collected from NW Wisconsin in May 1996 and examined for helminth parasites. Two species of helminths infected the spotted salamanders, while 3 species infected the red-backed salamanders. The nematode *Batracholandro magnavulvaris* had the highest prevalence in spotted salamanders (45%), while the nematode *Rhabdias* sp. had the highest prevalence in red-backed salamanders (30%). The trematode *Brachycoelium salamandrae* had the highest mean intensity in both hosts, 3.3 in *A. maculatum* and 2.0 in *P. c. cinereus*. This is the first report of *B. magnavulvaris* from *Ambystoma maculatum* as well as the first report of it from Wisconsin.

KEY WORDS: *Ambystoma maculatum*, *Plethodon c. cinereus*, *Brachycoelium salamandrae*, *Batracholandro magnavulvaris*, *Rhabdias* sp., Wisconsin.

The spotted salamander *Ambystoma maculatum* Shaw, 1802, is a large, robust species of mole salamander reported from south-central Ontario to Nova Scotia, south to Georgia and eastern Texas (Vogt, 1981). The red-backed salamander *Plethodon c. cinereus* Green, 1818, is one of the smallest woodland species of lungless salamanders that occurs throughout the northeastern United States and southeastern Canada, with populations in Ontario, Minnesota, Missouri, Arkansas, Oklahoma, Louisiana, and Georgia (Vogt, 1981). Both species inhabit mesic forests throughout northern Wisconsin. Although parasites of the spotted salamander, *Ambystoma maculatum*, and the red-backed salamander, *Plethodon c. cinereus*, have been studied by several

Table 1. Prevalence, abundance, and mean intensity of helminths of *Ambystoma maculatum* and *Plethodon c. cinereus*.

	<i>Ambystoma maculatum</i>			<i>Plethodon c. cinereus</i>			Location
	Prevalence*	Abundance ±1 SD	Mean intensity ±1 SD (range)	Prevalence*	Abundance ±1 SD	Mean intensity ±1 SD (range)	
<i>Brachycoelium salamandrae</i>	3 (15)	0.5 ± 1.3	3.3 ± 1.5 (2–5)	2 (10)	0.2 ± 0.6	2 (2)	Small and large intestine
<i>Batracholandros magnavulvaris</i>	9 (45)	0.8 ± 1.1	1.7 ± 1.1 (1–4)	1 (5)	0.05 ± 0.2	1 (1)	Small and large intestine
<i>Rhabdias</i> sp.	—	—	—	6 (30)	0.4 ± 0.8	1.3 ± 0.8 (1–3)	Body cavity

* Number infected (percent infected).

authors (Chitwood, 1933; Rankin, 1937a, b, 1938, 1945; Rankin and Hughes, 1937; Fischthal, 1955a, b; Cheng, 1958, 1960; Cheng and Chase, 1960; Ernst, 1974; Rosen and Manis, 1976; Dunbar and Moore, 1979; Bursey and Schibli, 1995), few studies are known from the Great Lakes area (Meserve, 1943; Coggins and Sajdak, 1982; Muzzall, 1990). Here we present new information on the parasites of Wisconsin salamanders.

Twenty adult red-backed salamanders (15 males and 5 females) were collected by overturning rocks and logs during the day, and 20 breeding spotted salamanders (15 males and 5 females) were collected by dip-net at an ephemeral pond in Bayfield County, Wisconsin, in May 1996. Animals were killed in MS 222 (ethyl m-aminobenzoate methane sulfonic acid) within 48 hours of capture. Snout-vent length and wet weight were recorded for each individual. The mean Snout-vent length ±1 SD (range) of spotted salamanders was 76 mm ± 8.8 (63–93). The mean Snout-vent length of red-backed salamanders was 44 mm ± 7 (30–52). At necropsy, the digestive tract, limb and body wall musculature, and internal organs were examined for helminths. Trematodes were preserved in 10% neutral buffered formalin, stained with aceto-carmine, dehydrated through ethanol, and mounted in Canada balsam. Nematodes were preserved in 10% neutral buffered formalin, dehydrated to 70% ethanol, cleared in glycerol, and identified as temporary mounts. Prevalence is the percentage of infected salamanders in a sample, mean intensity is the mean number of worms per infected salamander, and abundance is the mean number of individuals of a particular parasite species per host examined. Voucher specimens have been deposited in the Harold W. Manter

Laboratory, University of Nebraska State Museum, Lincoln (accession numbers HWML 39248—*Brachycoelium salamandrae* from red-backed salamander; 39249—*Brachycoelium salamandrae* from spotted salamander; 39250—*Rhabdias* sp.; 39251—*Batracholondros magnavulvaris*).

Eleven of 20 (55%) spotted salamanders were infected with helminths. Mean helminth abundance in spotted salamanders was 1.25 ± 1.60 . Nine (45%) were infected with *Batracholandros magnavulvaris* Rankin, 1937, while 3 (15%) were infected with *Brachycoelium salamandrae* Frolich, 1789. Only one spotted salamander (5%) was concurrently infected with *B. magnavulvaris* and *B. salamandrae*. Eight of 20 (40%) red-backed salamanders were infected with 1 or more *B. magnavulvaris*, *B. salamandrae*, and *Rhabdias* sp.. Mean helminth abundance in red-backed salamanders was 0.65 ± 0.99 . Only one red-backed salamander (5%) was concurrently infected with *B. magnavulvaris* and *B. salamandrae*. *Batracholandros magnavulvaris* had the highest prevalence in spotted salamanders, *Rhabdias* sp. had the highest prevalence in red-backed salamanders, and *B. salamandrae* had the highest mean intensity in both hosts (Table 1). There was no statistically significant correlation between helminth abundance and either weight or length for spotted salamanders ($r = -0.228$, $r = -0.111$) or red-backed salamanders ($r = 0.027$, $r = -0.096$). Because of the skewed sex ratio toward males, neither prevalence nor mean intensity was compared between the sexes in either host species.

Batracholandros magnavulvaris exhibits little host specificity, infecting plethodontids (Rankin, 1937b; Schad, 1960, 1963; Dyer et al., 1980; Goater et al., 1987; Muzzall, 1990; Joy et al.,

1993; McAllister et al., 1995; Bursey and Schibli, 1995) as well as salamandrids (Rankin, 1937b; Baker, 1987) and ambystomids (Muzzall and Schindlerle, 1992). In this study, 16 gravid female *B. magnavulvaris* were recovered from both salamander species, 15 from spotted salamanders and 1 from red-backed salamanders. All female nematode populations in salamanders has been reported by other investigators (Rankin, 1937b; Walton, 1940; Fischthal, 1955a; Joy et al., 1993). The spotted salamander is a new host record for *B. magnavulvaris*, and Wisconsin is a new locality record for this nematode in red-backed salamanders. Although both species of salamanders in this study are considered terrestrial, only the spotted salamander returns to water during its reproductive season. The red-backed salamander is totally terrestrial, with reproduction and development occurring on land (Vogt, 1981). Our prevalence data of *B. magnavulvaris* in spotted salamanders (45%) and red-backed salamanders (5%) agree with reports by Dunbar and Moore (1979) and Joy et al. (1993) that more aquatic salamanders are more likely to be infected with this nematode than are terrestrial species of salamanders. Although the differences observed in the present report may be due to chance, because of the small number of salamanders examined, Muzzall (1990) found 48 (28%) of 171 red-backed salamanders infected with this nematode, while Ernst (1974) found a 50% prevalence (6/12) in Virginia red-backed salamanders.

The 4 specimens of *Brachycoelium salamandrae* from red-backed salamanders and 10 specimens from spotted salamanders exhibited morphological variation. A statistically significant difference existed between the mean length, in millimeters \pm 1 SD (range), of trematodes recovered from spotted salamanders, 2.82 mm \pm 0.30 (2.28–3.30), and red-backed salamanders, 1.26 mm \pm 0.47 (0.85–1.93) (one-tailed *t*-test, *P* < 0.05), while no such difference existed in mean width of trematodes from spotted salamanders, 0.40 mm \pm 0.07 (0.28–0.50), and red-backed salamanders, 0.34 mm \pm 0.12 (0.25–0.52) (one-tailed *t*-test, *P* > 0.05). Other apparent differences existed in body shape, position of testes, and distribution of vitellaria. *Brachycoelium* spp. are common parasites of salamanders (Dyer et al., 1980; Goater et al., 1987; Muzzall, 1990; McAllister et al., 1995; Bursey and Schibli, 1995), but controversy surrounds the as-

signment of species to this genus. Rankin (1938) reduced all known species to *B. salamandrae*. He concluded that heavy infections produce many small flukes, whereas light infections were usually made up of larger specimens. Other investigators (Parker, 1941; Cheng, 1958, 1960; Cheng and Chase, 1960; Couch, 1966) disagreed and described 13 species based on such morphological characteristics as body length and shape, length of esophagus, position of testes, and distribution of vitellaria. Although the trematodes we collected from spotted salamanders and red-backed salamanders exhibited morphological variation, we have adopted the conservative approach suggested by McAllister et al. (1995) to report *B. salamandrae* from North American salamanders.

Rhabdias sp. was the most frequently found parasite in red-backed salamanders. Six red-backed salamanders (30%) were found to be infected, with a total of 8 specimens recovered from the body cavity. *Rhabdias* spp. have been reported previously from a wide variety of salamander species (Chitwood, 1933; Walton, 1938, 1940; Lehmann, 1954; Landewe, 1963; Dyer and Peck, 1975; Price and St. John, 1980; Coggins and Sajdak, 1982; Muzzall and Schindlerle, 1992). However, the finding of this nematode in the red-backed salamander was unexpected, since it is a lungless salamander and should not be expected to harbor lung parasites. The study of Baker (1979) on *Rhabdias* revealed that subadult nematodes must invade the lungs if they are to mature and produce eggs, but numerous subadults can be found in the body cavity. All nematodes recovered in the present study were nongravid subadults from the body cavity and are probably accidental infections in this host.

Results of the current survey support previous work on salamander helminths, indicating that they are not strongly host-specific; rather, their distribution can be correlated with habitat preference and diet of the host (Fischthal, 1955a, b; Dunbar and Moore, 1979; Coggins and Sajdak, 1982; Goater et al., 1987). Recently, Kleeberger and Werner (1982, 1983) showed that in a northern hardwood forest, similar to the present study area, spotted salamanders spent 72% of their time underground, 21% under decaying logs, and 7% under wet leaf litter, while red-backed salamanders in the same area spent 61% of their time above the soil, primarily in the litter layers

and decaying logs. Thus, the differences in parasite prevalence and species found in the two hosts may be due to differences in host habitat utilization. Parasites may encounter only a limited number of potential hosts under natural conditions, giving the appearance of a much narrower host specificity (Kennedy, 1975). To our knowledge, the spotted salamander is a new host record for *B. magnavulvaris*, and Wisconsin is a new locality record for this nematode in either of the salamander species.

We thank H. R. Yoder, University of Wisconsin–Milwaukee, and Dr. H. Gene Drecktrah, University of Wisconsin–Oshkosh, for help in collecting salamanders. We also thank two anonymous reviewers for improvements on an earlier draft of the manuscript.

Literature Cited

- Baker, M. R.** 1979. The free living and parasitic development of *Rhabdias* spp. (Nematoda: Rhabdiasidae) in amphibians. *Canadian Journal of Zoology* 57:161–167.
- . 1987. Synopsis of the Nematoda parasitic in amphibians and reptiles. *Memorial University of Newfoundland Occasional Papers in Biology* 11: 1–325.
- Bursey, C. R., and D. R. Schibli.** 1995. A comparison of the helminth fauna of two *Plethodon cinereus* populations. *Journal of the Helminthological Society of Washington* 62:232–236.
- Cheng, T. C.** 1958. Studies on the Trematode family Dicrocoeliidae. I. The genera *Brachycoelium* (Dujardin, 1845) and *Leptophallus* Luhe, 1909 (Brachycoeliinae). *American Midland Naturalist* 59: 67–81.
- . 1960. The life history of *Brachycoelium obseum* Nicoll, 1914, with a discussion of the systematic status of the trematode family Brachycoeliidae Johnston, 1912. *Journal of Parasitology* 46:464–474.
- , and **R. S. Chase, Jr.** 1960. *Brachycoelium stablefordi*, a new parasite of salamanders; and a case of abnormal polylobation of the testes of *Brachycoelium storeriae* Harwood, 1932. (Trematoda; Brachycoeliidae). *Transactions of the American Microscopical Society* 80:33–38.
- Chitwood, B. G.** 1933. On some nematodes of the superfamily Rhabditoidea and their status as parasites of reptiles and amphibians. *Journal of the Washington Academy of Science* 23:508–520.
- Coggins, J. R., and R. A. Sajdak.** 1982. A survey of helminth parasites in the salamanders and certain anurans from Wisconsin. *Proceedings of the Helminthological Society of Washington* 49:99–102.
- Couch, J. A.** 1966. *Brachycoelium ambystomae* sp. n. (Trematoda: Brachycoeliidae) from *Ambystoma opacum*. *Journal of Parasitology* 52:46–49.
- Dunbar, J. R., and J. D. Moore.** 1979. Correlations of host specificity with host habitat in helminths parasitizing the plethodontids of Washington County, Tennessee. *Journal of the Tennessee Academy of Science* 54:106–109.
- Dyer, W. G., and S. B. Peck.** 1975. Gastrointestinal parasites of the cave salamander, *Eurycea lucifuga* Rafinesque, from the southeastern United States. *Canadian Journal of Zoology* 53:52–54.
- , **R. A. Brandon, and R. L. Price.** 1980. Gastrointestinal helminths in relation to sex and age of *Desmognathus fuscus* (Green, 1818) from Illinois. *Proceedings of the Helminthological Society of Washington* 50:95–99.
- Ernst, E. M.** 1974. The parasites of the red-backed salamander, *Plethodon cinereus*. *Bulletin of the Maryland Herpetological Society* 10:108–114.
- Fischthal, J. H.** 1955a. Ecology of worm parasites in south-central New York salamanders. *American Midland Naturalist* 53:176–183.
- . 1955b. Helminths of salamanders from Promised Land State Forest Park, Pennsylvania. *Proceedings of the Helminthological Society of Washington* 22:46–48.
- Goater, T. M., G. W. Esch, and A. O. Bush.** 1987. Helminth parasites of sympatric salamanders: ecological concepts at infracommunity, component and compound community levels. *American Midland Naturalist* 118:289–300.
- Joy, J. E., T. K. Pauley, and M. L. Little.** 1993. Prevalence and intensity of *Thelandros magna-vulvaris* and *Omia papillocauda* (Nematoda) in two species of desmognathine salamanders from West Virginia. *Journal of the Helminthological Society of Washington* 60:93–95.
- Kennedy, C. R.** 1975. *Ecological animal parasitology*. John Wiley & Sons, Inc., New York. 531 pp.
- Kleeberger, S. R., and J. K. Werner.** 1982. Home range and homing behavior of *Plethodon cinereus* in Northern Michigan. *Copeia* 1982:409–415.
- , and ———. 1983. Post-breeding migration and summer movement of *Ambystoma maculatum*. *Journal of Herpetology* 17:176–177.
- Landewe, J. E.** 1963. Helminth and arthropod parasites of salamanders from southern Illinois. Unpublished M.S. Thesis, Southern Illinois University, Carbondale. 47 pp.
- Lehmann, D. L.** 1954. Some helminths of West Coast urodeles. *Journal of Parasitology* 40:231.
- McAllister, C. T., S. J. Upton, and S. E. Trauth.** 1995. Metazoan parasites of the graybelly salamander, *Eurycea multiplicata griseogaster* (Caudata: Plethodontidae), from Arkansas. *Journal of the Helminthological Society of Washington* 62: 70–73.
- Meserve, F. G.** 1943. *Phyllodistomum coatneyi* n. sp., a trematode from the urinary bladder of *Ambystoma maculatum* (Shaw). *The Journal of Parasitology* 29:226–228.
- Muzzall, P. M.** 1990. Endoparasites of the red-backed salamander, *Plethodon c. cinereus*, from southwestern Michigan. *Journal of the Helminthological Society of Washington* 57:165–167.
- , and **D. B. Schindlerle.** 1992. Helminths of the salamanders *Ambystoma t. tigrinum* and *Ambystoma t. tigrinum*.

- bystoma laterale* (Caudata: Ambystomatidae) from southern Michigan. Journal of the Helminthological Society of Washington 59:201–205.
- Parker, M. V.** 1941. The trematode parasites from a collection of amphibians and reptiles. Journal of the Tennessee Academy of Science 16:27–44.
- Price, R. L., and T. St. John.** 1980. Helminth parasites of the small-mouth salamander, *Ambystoma texanum* Matthes, 1855, from William County, Illinois. Proceedings of the Helminthological Society of Washington 47:273–274.
- Rankin, J. S.** 1937a. An ecological study of parasites of some North Carolina salamanders. Ecological Monographs 7:171–269.
- . 1937b. New helminths from North Carolina salamanders. Journal of Parasitology 23:29–42.
- . 1938. Studies on the trematodes genus *Brachycoelium* Duj. I. Variation in specific characters with reference to the validity of the described species. Transactions of the American Microscopical Society 57:358–375.
- . 1945. An ecological study of the helminth parasites of amphibians and reptiles of western Massachusetts and vicinity. Journal of Parasitology 31:142–152.
- , and **R. C. Hughes.** 1937. Notes on *Diplostomulum ambystomae* n. sp. Transactions of the American Microscopical Society 57:61–66.
- Rosen, R., and R. Manis.** 1976. Trematodes of Arkansas amphibians. Journal of Parasitology 62: 833–834.
- Schad, G. A.** 1960. The genus *Thelandros* (Nematoda: Oxyuroidea) in North American salamanders, including a description of *Thelandros salamandrae* n. sp. Canadian Journal of Zoology 38: 115–120.
- . 1963. *Thelandros magnavulvaris* (Rankin, 1937) Schad, 1960 (Nematoda: Oxyuroidea) from the green salamander, *Aneides aeneus*. Canadian Journal of Zoology 41:943–946.
- Vogt, R. C.** 1981. Natural History of Amphibians and Reptiles of Wisconsin. Milwaukee Public Museum and Friends of the Museum, Milwaukee. 205 pp.
- Walton, A. C.** 1938. The Nematoda as parasites of Amphibia IV. Transactions of the American Microscopical Society 57:38–53.
- . 1940. Some nematodes from Tennessee amphibia. Journal of the Tennessee Academy of Science 15:402–405.

J. Helminthol. Soc. Wash.
65(1), 1998 pp. 102–104

Research Note

Nematodes of the Great Plains Narrow-mouthed Toad, *Gastrophryne olivacea* (Microhylidae), from Southern Arizona

STEPHEN R. GOLDBERG,¹ CHARLES R. BURSEY,² AND HAY CHEAM¹

¹ Department of Biology, Whittier College, Whittier, California 90608 (e-mail: sgoldberg@whittier.edu)

² Department of Biology, Pennsylvania State University, Shenango Campus, 147 Shenango Avenue, Sharon, Pennsylvania 16146 (e-mail: cxb13@psuvm.psu.edu)

ABSTRACT: Thirty *Gastrophryne olivacea* from southern Arizona were examined for helminths. Two species of nematodes, *Aplectana incerta* and *Aplectana itzocanensis*, were found. Both represent new host records. *Aplectana itzocanensis* had the higher prevalence (70%) and greater mean intensity (13.0). In southern Arizona, *A. incerta* and *A. itzocanensis* also occur concurrently in other anurans.

KEY WORDS: *Gastrophryne olivacea*, Microhylidae, Nematodes, *Aplectana incerta*, *Aplectana itzocanensis*, Arizona.

The Great Plains narrowmouth toad, *Gastrophryne olivacea* (Hallowell, 1856), occurs from eastern Nebraska and western Missouri, through Oklahoma and Texas, west through northern México, into south central Arizona, from sea

level to 1250 m elevations (Stebbins, 1985). There are several reports of helminths in *G. olivacea*: Kansas (Freiburg, 1951), Oklahoma (Kuntz, 1941), and Texas (Harwood, 1932; McAllister and Upton, 1987). The purpose of this note is to report nematodes from a population of *G. olivacea* from southern Arizona.

Thirty *G. olivacea* (8 females, 22 males, mean snout-vent length = 27 mm ± 2.3 SD; range, 23–32 mm) were borrowed from the herpetology collection of the University of Arizona, Tucson (UAZ) 15824, 15826, 15827, 15830, 15833, 20561, 20563, 25858, 25861, 25863, 29026, 29027, 29032, 29035, 29036, 29039–29041, 29043, 29046–29049, 29051, 29052, 29055–29058, 32016. These specimens had been col-